



## Continuous monitoring and modelling of cyanobacteria dynamics in urban lakes

Talita Silva, Brigitte Vinçon-Leite, Bruno J. Lemaire, Bruno Tassin, Martin Seidl, Nilo Nascimento

### ► To cite this version:

Talita Silva, Brigitte Vinçon-Leite, Bruno J. Lemaire, Bruno Tassin, Martin Seidl, et al.. Continuous monitoring and modelling of cyanobacteria dynamics in urban lakes. 2nd IWA Symposium on Lake and Reservoir Management, Jun 2011, Grenade, Spain. hal-00674661


HAL Id: hal-00674661

<https://enpc.hal.science/hal-00674661>

Submitted on 28 Feb 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



École des Ponts  
ParisTech



UFMG



# Continuous monitoring and modelling of cyanobacteria dynamics in urban lakes

Application to Lake Enghien, France and Lake Pampulha, Brazil

**Talita Silva, Brigitte Vinçon-Leite, Bruno Lemaire, Bruno Tassin,  
Martin Seidl, Nilo Nascimento**



# OUTLINE

- Cyanobacteria monitoring
- Proliphyc project
  - Measurement buoy
- Lake Enghien
  - Monitoring
  - Modelling
- Lake Pampulha
  - Monitoring methodology
  - Integrated modelling approach
  - Expected results
- Conclusions

# LIMITS OF THE USUAL MONITORING

- Cyanobacteria monitoring



- How it is usually done




Leesu (2010)



Cyanobacteria – Chlorophyll-a concentration ( $\mu\text{g. L}^{-1}$ )





- How we propose to do it



# BENEFITS OF HIGH-FREQUENCY MONITORING

## French protocol (European WFD)



- 4 sampling campaigns/year
  - Including 3 during the summer
- 3 weeks = minimum interval between the campaigns



4


# PROLIPHYC PROJECT

- A measuring buoy
  - High-frequency data (15-30 min)
  - Data teletransmission (daily)
- Warning system
  - Based on the weather forecast and buoy measurements
  - Short-term predictive modelling (1 week)




# LAKE ENGHIEN-LES-BAINS

- 41 ha, mean depth = 1.5 m
- Rainwater storing, sport, landscape
- Inappropriate wastewater discharges
- *Planktothrix agardhii* blooms




## MODELLING

- DYRESM-CAEDYM model (Centre for Water Research, University of Western Australia)
- 1 phytoplankton group: cyanobacterium *Planktothrix agardhii*
- Model calibration and validation



# DYCD RESULTS

- Model results show good agreement with measurements
- High frequency data → thorough assessment of results
  - Gal *et al.* (2009); Burger *et al.* (2008); Hornung (2002)
  - Medium-term simulations, hourly time step




## LAKE PAMPULHA

- Belo Horizonte, Minas Gerais, Brazil
- 197 ha, 10 hm<sup>3</sup>, 5,0 m (mean depth)
- Former drinking water source

Urbanization + no sanitation  
infrastructure + no erosion control

lake silting and storage capacity  
reduction

- Cyanobacteria blooms and  
macrophyte growth
- Touristic spot
  - Fishing, sailing, landscape



# WATERSHED AND LAKE MONITORING

- High frequency data\*

- Two main tributaries and lake outlet :
  - Rain gauges
  - Flow rate
  - Temperature, conductivity and turbidity
  - Automatic samplers: nutrients and dissolved oxygen
- Measurement buoy :
  - Temperature, conductivity, pH, O<sub>2</sub> and chl-a
- Meteorological forcing : INMET




10

\*MAPLU project: Funded by FINEP (Brazilian Research Agency)

# INTEGRATED APPROACH


- Watershed hydrological model +Lake ecological model



Why a modelling approach?

## INTEGRATED APPROACH




- Watershed hydrologic model +Lake ecological model



Lake response x Different scenarios of watershed evolution:  
Land-use changes, impervious area expanding...

## LAKE ECOLOGICAL MODEL

- Belo Horizonte municipality data
- Cyanobacteria species:
  - *Nostocaceae, Microcystis wesenbergii, Sphaerocavum sp.*
- Cyanobacteria modelling: on-going



# CONCLUSIONS

- Lake Enghien
  - Warning system
  - High frequency lake monitoring
  - Cyanobacteria dynamics modelling
  - Few data about the catchment
- Lake Pampulha
  - Management tool
  - High frequency watershed monitoring *in progress*
  - High frequency lake monitoring *from September/2011*
  - Cyanobacteria dynamics modelling *preliminary results*
  - Watershed modelling and scenarios

NEWSBULLETIN AN OCCURRENCE OF CYANOBACTERIA (BLUE-GREEN ALGAE)  
HAS BEEN DETECTED IN CHAMCOOK LAKE



**THANK YOU FOR YOUR ATTENTION!**