

Eastward-moving Equatorial Modons in moist-convective Shallow-Water Models

Masoud Rostami, Vladimir Zeitlin

▶ To cite this version:

Masoud Rostami, Vladimir Zeitlin. Eastward-moving Equatorial Modons in moist-convective Shallow-Water Models. Geophysical and Astrophysical Fluid Dynamics, 2020, pp.1-23. 10.1080/03091929.2020.1805448 . hal-02919418

HAL Id: hal-02919418 https://enpc.hal.science/hal-02919418

Submitted on 22 Aug 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Eastward-moving Equatorial Modons in moist-convective Shallow-Water Models

MASOUD ROSTAMI^{a,b}, VLADIMIR ZEITLIN^a

 $^a {\rm Laboratoire}$ de Météorologie Dynamique (LMD)/IPSL, Sorbonne Université and Ecole Normale Supérieure (ENS), Paris, France

 b Institute for Geophysics and Meteorology (IGM), University of Cologne, Cologne, Germany

This is the Author's Original Manuscript (AOM); that is, the manuscript in its original form; a "preprint". The Version of Record of this manuscript has been accepted for publication by the Journal of *Geophysical & Astrophysical Fluid Dynamics*, published by *Taylor & Francis*. The publisher's version is available at https://doi.org/10.1080/03091929.2020.1805448.

Geophysical and Astrophysical Fluid Dynamics Vol. 00, No. 00, 00 Month 20xx, 2–22

Eastward-moving Equatorial Modons in moist-convective Shallow-Water Models.

MASOUD ROSTAMI^{\dagger} [‡] and VLADIMIR ZEITLIN [†]

[†] Laboratoire de Météorologie Dynamique/Sorbonne Université/ Ecole Normale Supérieure (ENS)/CNRS, 24 Rue Lhomond, 75005 Paris, France

[‡] Institute for Geophysics and Meteorology, University of Cologne, 50969 Cologne, Germany (Received 00 Month 20xx; final version received 00 Month 20xx)

It is shown that steady large-scale slowly eastward-moving twin-cyclone coherent 1 structures, the equatorial modons, exist in both one- and two layer versions of the 2 rotating shallow water model on the equatorial beta plane. They arise via the pro-3 cess of "ageostrophic adjustment" from the analytic asymptotic modon solutions of 4 the vorticity equation obtained in the limit of small pressure perturbations. Evolu-5 tion of these structures in adiabatic and moist-convective environments, and also in 6 the presence of topography is analyzed, showing their robustness in the one-layer 7 model. It is demonstrated that moist convection enhances and helps maintain the 8 modons. In the two-layer model the barotropic and quasi-barotropic modons display 9 similar to one-layer modon features, while increasing baroclinicity leads to eventual 10 loss of coherence and arrest of the eastward propagation. Some features of equa-11 torial modons resemble those observed in the Madden-Julian Oscillation events in 12 tropical atmosphere, which hints at their possible relevance to the dynamics of this 13 phenomenon. 14

Keywords: Equatorial Waves, Madden Julian Oscillation, Modon, Moist Convection, Rotating Shallow 15 Water Model 16

1. Introduction 17

Atmospheric dynamics in the equatorial regions of rotating planets significantly differs from 18 that in higher latitudes due to the simple fact that the rotation axis lies in the tangent plane 19 to the equator. As is well known starting from the pioneering paper by Matsuno (1966), 20 equatorial dynamics is dominated by specific equatorial waves, e.g. (Gill 1982). These waves 21 are well identified in meteorological in situ and satellite observations, e.g. (Wheeler and Kiladis 22 1999). Yet, both kinds of observations reveal slow eastward-propagating structures which do 23 not fall in any class of equatorial waves. They were identified as periodically arising large-scale 24 patterns of enhanced deep convection which are slowly moving from the Indian Ocean over the 25 maritime continent and dying out in the Pacific, the Madden-Julian Oscillation (MJO). After 26 its discovery (Madden and Julian 1972), much was learnt about its structure and properties, 27 e.g. Zhang (2005). Yet dynamical nature of MJO is not fully understood (Zhang 2005) and, 28 in particular, there is no clear answer to the question why MJO events are slowly moving 29 eastward. Climate models have difficulties in reproducing them, e.g. (Kim et al. 2011). The 30 Kelvin waves, the only species of the equatorial waves that can produce steady large-scale 31

Equatorial Modons

³² eastward-moving structures, have much faster velocity than MJO (Straub and Kiladis 2002).

³³ There exist simplified models of MJO, either quasi-linear, e.g. Majda and Stechman (2009),

Sobel and Maloney (2012), or nonlinear Haertel et al. (2013), all of them heavily relying on the dynamical role of moist convection.

The question however arises, whether there exist "dry" dynamical structures moving slowly 36 eastwards along the equator, which could provide a dynamical "backbone" of the MJO events. 37 The spectrum of linear equatorial waves being well established, such objects should be nec-38 essarily nonlinear. There do exist steady eastward-moving structures in a fluid layer in the 39 mid-latitude tangent- plane approximation, the so-called modons, exact dipolar solutions of 40 the quasi-geostrophic (QG) equations discovered by Larichev and Reznik (1976). Although 41 the classical modons were found analytically in the QG approximation, it was shown first by 42 a computer-assisted analysis (Kizner et al. 2008) and then by direct numerical simulations 43 (Ribstein et al. 2010) that corresponding solutions exist also in the parent rotating shallow 44 water (RSW) model on the f- plane. The QG modon solutions were extended to the full 45 sphere (Verkley 1984, Tribbia 1984) in the framework of equivalent barotropic model, which 46 is structurally close to QG. In their pioneering paper, Yano and Tribbia (2017) came up with 47 an idea that the MJO could be related to such a modon in spherical geometry and showed 48 that, like on the tangent plane (Ribstein et al. 2010), the equivalent barotropic modon persists 49 in the parent RSW model on the sphere. Yet, the archetype model for understanding dynam-50 ics of the tropical atmosphere is RSW on the equatorial beta-plane, e.g. Matsuno (1966). For 51 example, the spectrum of convectively coupled equatorial waves, including MJO, is studied by 52 projection onto the wave-spectrum of the RSW model (Wheeler and Kiladis 1999, Straub and 53 Kiladis 2002). The idea, thus, arises to look for modons in this model. However, it encounters 54 an obstacle from the very beginning because a consistent QG approximation at the equator 55 is problematic, e.g. James (1994), so the known QG modon solutions can not be directly bor-56 rowed. We should emphasize at this point, that coherent westward-propagating dipoles were 57 found in the RSW model on the equatorial beta-plane by Boyd (1985), and were also called 58 modons. However, their nature is different, as they correspond to the KdV solitons of weakly 59 nonlinear long equatorial Rossby waves, see (LeSommer et al. 2004). A conjecture that east-60 ward propagating modons could exist was also formulated by Boyd (1985). We will discuss 61 this point in Sect. 3 below. 62

Let us also emphasize that the applications of the RSW model on the equatorial beta-plane are not limited to tropical meteorology. The related equatorial wave dynamics is regularly invoked for explanations of the observations in the equatorial regions of planetary atmospheres, e.g. (Showman and Dowling 2000, Legarreta et al. 2016, Sanchez-Lavega et al. 2017).

Our main observation (Rostami and Zeitlin 2019a) is that there exist a dynamical regime 67 in the RSW model, corresponding to the so-called long-wave approximation in oceanography 68 (Gill 1982), which arises in the limit of small pressure variations and gives, to the leading or-69 der, a vorticity equation which does allow for modon solutions. This quasi-barotropic regime is 70 the RSW analog of Charney (1963) non-divergent balance model for equatorial atmosphere. 71 The relevance of this model to large-scale tropical motions is supported by scale and data 72 analyses (Yano and Bonazzola 2009, Yano et al. 2009). Modon solutions in this regime were 73 constructed in Rostami and Zeitlin (2019a). Following the strategy of Ribstein et al. (2010), 74 these asymptotic solutions were then used to initialize high-resolution numerical simulations 75 with the full RSW model, and to show that coherent dipolar steady eastward-moving struc-76 tures, the RSW modons, do arise and persist. However, the domain of attraction of such 77 modon solutions seems to be limited as they do not arise without added dipolar vorticity 78 component from initial localized circular pressure anomalies, even if these latter have consid-79 erable amplitudes. The modon configurations, however do arise from such configurations in 80 the presence of moist-convective feedback (Rostami and Zeitlin 2019b). 81

⁸² Below we recall the derivation of equatorial modon solutions in one-layer RSW and present

M. ROSTAMI & V. ZEITLIN

numerical simulations illustrating the influence of the effects of moisture and topography on 83 their behavior. We show that topographic obstacles of reasonable amplitude do not disrupt 84 the modons, and that inclusion of moisture, with condensation and evaporation, enhances 85 the modons, and produces some specific convective and condensation patterns. We then turn 86 87 to baroclinic generalizations of one-layer modons, and look for similar solutions in the twolayer equatorial RSW model. We analyze different parameter regimes in this model, under the 88 same hypothesis of small pressure variations, and explore the possibilities of having coherent 89 dipolar solutions. We show that such asymptotic solutions do exist, verifying layer-wise in 90 the leading order the same vorticity equation as in the one-layer RSW. We then follow the 91 same strategy as in the one-layer case, and initialize high-resolution numerical simulations 92 with well-balanced finite-volume scheme for two-layer RSW equations on the equatorial beta-93 plane with the velocity fields corresponding to such asymptotic solutions. The simulations 94 reveal that only close to barotropic initializations, i.e. those with the same in each layer, 95 or close to, initial asymptotic modons, give rise to slowly eastward-moving two-layer RSW 96 modons, while baroclinic, i.e. with asymptotic modons of essentially different size and/or 97 intensity, initializations result in dipolar structures which are eventually turning westward 98 and becoming equatorial Rossby-wave packets. 99

The paper is organized as follows. We first recall the theory of barotropic equatorial modons in one-layer equatorial RSW, and then analyze its generalization to the two-layer case in Sect.

2. We present the results of numerical simulations initialized with asymptotic modon solutions
in both one- and two-layer cases, and in both "dry" and moist-convective environments in

¹⁰⁴ Sect. 3. Section 4 contains summary and discussion.

105 2. Modons in one- and two-layer RSW models: theory

106 2.1. One-layer configuration

107 2.1.1. From one-layer RSW to barotropic model

In this Section we recall the construction of equatorial modons which was sketched in 108 Rostami and Zeitlin (2019a). The RSW model we are using should be understood as vertically 109 averaged primitive equations in pseudo-height pressure coordinates (Zeitlin 2018), and can be 110 extended to include water vapor with condensation and related latent heat release (Bouchut 111 et al. 2009), as well as surface evaporation (Lahaye and Zeitlin 2016, Rostami and Zeitlin 2017). 112 Liquid water, precipitation, vaporisation and related cooling can be also incorporated into the 113 model, as well as vertical structure, see Rostami and Zeitlin (2018) for these generalizations, 114 and references therein for previous successful applications of the model. Below we recall the 115 approximations leading to the Charney regime in one-layer RSW model on the equatorial 116 beta-plane following Rostami and Zeitlin (2019a). 117

¹¹⁸ The "dry" one-layer RSW equations in the equatorial beta- plane with no dissipation read:

$$\partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v} + \beta y \, \hat{\mathbf{z}} \wedge \mathbf{v} + g \nabla h = 0 \,, \tag{1}$$

119 120

121

$$\partial_t h + \nabla \cdot [\mathbf{v}(h-b)] = 0, \qquad (2)$$

where $\nabla = (\partial_x, \partial_y)$, $\mathbf{v} = (u, v)$, u is zonal and v- meridional components of velocity, h is thickness in pseudo-height coordinates, b > 0 represents topography, β is the meridional gradient of the Coriolis parameter, and $\hat{\mathbf{z}}$ is a unit vertical vector. We introduce a nonlinearity parameter λ , which measures the amplitude of pressure variations $h = H(1 + \lambda \eta)$, where His the unperturbed thickness, and rewrite the momentum equations correspondingly:

$$\partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v} + \beta y \, \hat{\mathbf{z}} \wedge \mathbf{v} + g H \lambda \nabla \eta = 0 \,, \tag{3}$$

Equatorial Modons

where we omitted topography, which would be reintroduced later. We then choose the same

where we omitted topography, which would be reintroduced later. We then choose the same spatial and velocity scales in both zonal and meridional directions $(x, y) \sim L$, $(u, v) \sim V$, and choose the time-scale as: $t \sim (L/V)$. Notice that such uniform in zonal and meridional directions scaling differs from the traditionally used one, cf. Gill (1980), LeSommer et al. (2004). The non-dimensional momentum equations are:

$$\partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v} + \overline{\beta} y \, \hat{\mathbf{z}} \wedge \mathbf{v} + \frac{g H \lambda}{V^2} \nabla \eta = 0 \,, \tag{4}$$

where we introduced a non-dimensional beta parameter $\overline{\beta} = \beta L^2 / V$, which is, in fact, the inverse equatorial Rossby number. We should emphasize that, unlike the traditional equatorial scaling, which uses the equatorial deformation radius $L_d = (gH)^{\frac{1}{4}} / \beta^{\frac{1}{2}}$, the spatial scale is not rigidly related to other parameters here. If we suppose that

$$\lambda \to 0, \text{ and } \frac{gH\lambda}{V^2} = \mathcal{O}(1),$$
(5)

which immediately gives $V \ll \sqrt{gH}$, that is the characteristic velocity is much smaller than the phase velocity of the fastest waves $c = \sqrt{gH}$, like Kelvin wave, then the momentum and mass conservation equations become:

138

133

$$\partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v} + \overline{\beta} y \, \hat{\mathbf{z}} \wedge \mathbf{v} + \nabla \eta = 0 \,, \tag{6}$$

$$\lambda(\partial_t \eta + \mathbf{v} \cdot \nabla \eta) + (1 + \lambda \eta) \nabla \cdot \mathbf{v} = 0.$$
⁽⁷⁾

In the leading order of the expansion in
$$\lambda$$
: $\mathbf{v} = \mathbf{v}_0 + \lambda \mathbf{v}_1 + \dots$ equation (7) gives $\nabla \cdot \mathbf{v}_0 = 0$, and the motion is divergenceless. Hence, we can introduce a streamfunction for the leading-order

the motion is divergenceless. Hence, we can introduce a streamfunction for the leading-order velocity:

$$u_0 = -\partial_y \psi, \, v_0 = \partial_x \psi. \tag{8}$$

First-order corrections u_1 , v_1 can be found in the next order of the asymptotic expansion in λ . There is no guarantee that such expansion converges, but numerical results presented below suggest that there is indeed a correspondence between solutions of the full equations (1), (2) and the asymptotic ones. Cross-differentiation of the zonal and meridional momentum equations allows to eliminate η and gives the vorticity equation for streamfunction:

$$\nabla^2 \psi_t + \mathcal{J}(\psi, \nabla^2 \psi) + \overline{\beta} \psi_x = 0, \tag{9}$$

153 where $\mathcal J$ denotes the Jacobian.

If topography would be of the order one, and would be kept, we would have $(H - b)u_0 = -\partial_y \psi$, $(H - b)v_0 = \partial_x \psi$ instead of (8), and the resulting system would become an equatorial version of the long-wave approximation, which is used for studies of the dynamical effects of topography in oceanography, cf. (Gill 1982). Here we will use it first without topography, in order to construct the modon solutions.

159 2.1.2. One-layer RSW modons

Equation (9) coincides with the infinite deformation radius limit of one-layer QG equa-160 tion, e.g. Zeitlin (2018), so the construction of the modon solutions can be straightforwardly 161 transposed, following the pioneering paper of Larichev and Reznik (1976). Let us recall that 162 these solutions are obtained under hypothesis of steady motion with constant zonal velocity 163 U, by supposing a linear relationship between the absolute vorticity and streamfunction in 164 the co-moving frame, which leads to an inhomogeneous Helmholtz equation. It is solved by 165 separation of variables in polar coordinates in terms of Bessel functions, first in the outer 166 with respect to a separatrix domain, under condition of decay at infinity, and then in the 167 inner domain, and matching the inner and outer solutions at the separatrix. The separatrix 168 is a circle of radius a in the (x, y) - plane. The matching conditions are continuity of the 169

172

M. ROSTAMI & V. ZEITLIN



Figure 1. Streamlines and velocity field of the asymptotic modon corresponding to the first root (a, p) = (0.5, 3.1623) of the equation (11), with slow eastward propagation U = 0.1, in stationary (*left panel*), and co-moving frames (*middle panel*), and a solution corresponding to the second root in the stationary frame (*right panel*). Dashed line: separatrix of radius a.

streamfunction and its normal derivative across the circle. It is worth noting that the actual
size of the vortex, as a nonlinear solution, is larger than a. The solution is:

$$\begin{cases} \psi_{ext} = -\frac{Ua}{K_1(pa)} K_1(pr) \sin \theta, \ r > a, \\ \\ \psi_{int} = \left[\frac{Up^2}{k^2 J_1(ka)} J_1(kr) - \frac{r}{k^2} (1 + U + Uk^2) \right] \sin \theta, \ r < a, \end{cases}$$
(10)

where J_1 and K_1 are ordinary and modified Bessel functions, respectively, p is real, and $p^2 = \overline{\beta}/U$, so U > 0, and the motion is *eastward*. The matching conditions give:

$$\frac{J_1'(ka)}{J_1(ka)} = \frac{1}{ka} (1 + \frac{k^2}{p^2}) - \frac{k}{p} \frac{K_1'(pa)}{K_1(pa)},\tag{11}$$

and for each pair (a, p) (or (a, U)) there exist a series of eigenvalues of k arising from (11), of 175 which the lowest corresponds to a vortex dipole, to be called the asymptotic modon. We will 176 be using below U as a governing parameter. The higher eigenvalues correspond to multi-polar 177 vortices with significant velocity shear inside the core, which are more sensitive to dissipation, 178 and are rarely considered in the literature. By construction, thus obtained solutions can be 179 centered anywhere in the equatorial beta-plane. Yet it is only when the center of the circle is 180 at the equator that the modon is a pair of cyclones, due to the change of sign of the planetary 181 vorticity βy , otherwise it is a cyclone-anticyclone pair. The streamlines of the asymptotic 182 modons corresponding to the first and the second roots of the equation (11) are presented in 183 Fig. 1. Pressure distribution at a given ψ is obtained by differentiating and combining the 184 leading-order momentum equations: 185

$$\nabla^2 \eta = Hess[\psi] + \overline{\beta}(\psi_y + y\nabla^2\psi) \tag{12}$$

where $Hess[\psi] = \partial_{xx}^2 \psi \, \partial_{yy}^2 \psi - (\partial_{xy}^2 \psi)^2$ is the Hessian of ψ . Notice that equations (9) and (12) are shallow-water counterparts of the non-divergent model for large-scale tropical motions derived in Charney (1963).

189 2.2. Two-layer configuration

190 2.2.1. The model, and small pressure perturbation limit(s)

¹⁹¹ We start with the "dry" two-layer atmospheric RSW momentum and mass equations in ¹⁹² the equatorial β -plane with no topography and dissipation. The model is sketched in Figure Equatorial Modons



Figure 2. A sketch of the two-layer atmospheric rotating shallow water model.

¹⁹³ 2 The equations of the model read:

194

$$\begin{cases} \partial_t \mathbf{v_1} + \mathbf{v_1} \cdot \nabla \mathbf{v_1} + \beta \, y \, \hat{\mathbf{z}} \wedge \mathbf{v_1} + g \nabla (h_1 + h_2) = 0 \,, \\\\ \partial_t \mathbf{v_2} + \mathbf{v_2} \cdot \nabla \mathbf{v_2} + \beta \, y \, \hat{\mathbf{z}} \wedge \mathbf{v_2} + g \nabla (h_1 + sh_2) = 0 \,, \\\\ \partial_t h_1 + \nabla \cdot (\mathbf{v_1}h_1) = 0 \,, \\\\ \partial_t h_2 + \nabla \cdot (\mathbf{v_2}h_2) = 0 \,, \end{cases}$$
(13)

where $s = \theta_2/\theta_1 > 1$ is a stratification parameter, $\theta_{1,2}$ are potential temperatures of the layers, and β is the meridional gradient of the Coriolis parameter, as before. We introduce the deviations η_1 and η_2 of the free surface and interface between the layers from their positions at rest, respectively, and corresponding nonlinearity parameters $\lambda_{1,2}$, as follows:

$$\begin{cases}
h_2 = H_2 - \eta_2 = H(d_2 - \lambda_2 \overline{\eta}_2), \\
h_1 = H_1 - \eta_1 + \eta_2 = H(d_1 - \lambda_1 \overline{\eta}_1 + \lambda_2 \overline{\eta}_2),
\end{cases}$$
(14)

Here $d_2 = H_2/H$, and $d_1 = H_1/H$ are non-dimensional thicknesses of the layers. d_1 and $d_2 = 1 - d_1$ are not independent. We nonetheless use both d_1 and d_2 explicitly in the formulas below to keep their expression concise. With the same choice of scaling as in the one-layer model for both layers, the non-dimensional equations (13) are:

$$\partial_{t} \mathbf{v}_{1} + \mathbf{v}_{1} \cdot \nabla \mathbf{v}_{1} + \overline{\beta} \, y \, \hat{\mathbf{z}} \wedge \mathbf{v}_{1} = -\frac{gH \lambda_{1}}{V^{2}} \nabla \overline{\eta}_{1},$$

$$\partial_{t} \mathbf{v}_{2} + \mathbf{v}_{2} \cdot \nabla \mathbf{v}_{2} + \overline{\beta} \, y \, \hat{\mathbf{z}} \wedge \mathbf{v}_{2} = -\frac{gH}{V^{2}} [\lambda_{1} \nabla \overline{\eta}_{1} - (1 - s)\lambda_{2} \nabla \overline{\eta}_{2}],$$

$$d_{1} \nabla \cdot \mathbf{v}_{1} - \lambda_{1} [\partial_{t} \overline{\eta}_{1} + \nabla (\overline{\eta}_{1} \mathbf{v}_{1})] + \lambda_{2} [\partial_{t} \overline{\eta}_{2} + \nabla (\overline{\eta}_{2} \mathbf{v}_{1})] = 0,$$

$$d_{2} \nabla \cdot \mathbf{v}_{2} - \lambda_{2} [\partial_{t} \overline{\eta}_{2} + \nabla (\overline{\eta}_{2} \mathbf{v}_{2})] = 0,$$
(15)

203

Even if we suppose that parameters s and $d_{1,2}$ which control the stratification are of the order one, there are still two small parameters $\lambda_{1,2}$ in the small pressure perturbations limit, and their ratio should be prescribed in order to fix a dynamical regime. There are two obvious choices: both λ_1 and λ_2 are of the same order, or one of them is much smaller than the other one. We will call them, respectively, Regime 1 and Regime 2, the second one having two sub-regimes depending on which of $\lambda_{1,2}$ is smaller.

M. ROSTAMI & V. ZEITLIN

210 2.2.2. Regime 1

In this regime deviations of the free surface and of the interface are of the same order: $\lambda_1 \sim \lambda_2 \sim \lambda \rightarrow 0$, hence $[(gH\lambda)/V^2] \sim 1$ and from the third and forth equations in (15) we get, analogously to the one-layer case: $\nabla \cdot \mathbf{v}_1 = \nabla \cdot \mathbf{v}_2 = 0$. Therefore,

214
$$\nabla^2 \psi_{(1,2)t} + \mathcal{J}(\psi_{(1,2)}, \nabla^2 \psi_{(1,2)}) + \overline{\beta} \psi_{(1,2)x} = 0, \tag{16}$$

where $\psi_{(1,2)}$ are streamfunctions in respective layers. Each of these equations has modon 215 solutions, but they are not necessarily the same and may differ in scale and/or intensity. 216 To the leading order, thus, the velocities in the layers do not influence each other, but the 217 pressures do. Indeed, by taking divergence of the momentum equation for layer 1, and using 218 the fact that to the leading order the divergence of velocity is zero, we get a diagnostic 219 equation for η_1 analogous to (12), which allows to determine η_1 in terms of velocity in layer 220 1. Similar manipulations with the momentum equation for layer 2 give a diagnostic equation 221 for η_2 which, however, contains η_1 , and thus is taking into account a coupling between the 222 layers. We should emphasize that the present dynamical regime is fully consistent with the 223 construction of Charney (1963) for continuously stratified atmosphere, where the horizontal 224 velocity obeying the equation (9) could have arbitrary modulation in the vertical. 225

226 2.2.3. Regime 2a

229

232

In this regime the deviation of the free surface is much smaller than that of the interface: $\lambda_1 \sim \lambda^2$, $\lambda_2 \sim \lambda \to 0$. Still supposing that $[(gH\lambda)/V^2] \sim 1$, we therefore have to rescale the lower-layer velocity: $\mathbf{v}_1 = \lambda \mathbf{v}_1^*$, and thus get:

$$\partial_t \mathbf{v}_1^* + \lambda \mathbf{v}_1^* \cdot \nabla \mathbf{v}_1^* + \overline{\beta} \, y \, \hat{\mathbf{z}} \wedge \mathbf{v}_1^* = -\nabla \overline{\eta}_1, \tag{17a}$$

$$\partial_t \mathbf{v_2} + \mathbf{v_2} \cdot \nabla \mathbf{v}_2 + \overline{\beta} \, y \, \hat{\mathbf{z}} \wedge \mathbf{v}_2 = (1 - s) \nabla \overline{\eta}_2, \tag{17b}$$

$$d_1 \nabla \cdot \mathbf{v}_1^* + \partial_t \overline{\eta}_2 = 0, \qquad (17c)$$

$$\nabla \cdot \mathbf{v}_2 = 0, \tag{17d}$$

Cross-differentiation and differentiation of (17a) give, respectively, to the leading order n λ :

$$\partial_t \zeta_1^* + \overline{\beta} \, y \, \nabla \cdot \mathbf{v}_1^* + \overline{\beta} \, v_1^* = 0, \tag{18a}$$

$$\partial_t (\nabla \cdot \mathbf{v}_1^*) - \overline{\beta} \, y \, \zeta_1^* + \overline{\beta} \, u_1^* = \nabla^2 \overline{\eta}_1, \tag{18b}$$

We introduce a potential and a streamfunction for the lower-layer velocity, which is not divergenceless, contrary to Regime 1:

$$\begin{cases} u_1^* = \phi_{1x}^* - \psi_{1y}^*, \\ v_1^* = \phi_{1y}^* - \psi_{1x}^*, \end{cases}$$
(19)

such that $\zeta_1^* = \nabla^2 \psi_1^*$ and $\nabla \cdot \mathbf{v}_1^* = \nabla^2 \phi_1^*$. Consequently Eq.(17c) can be rewritten as $\nabla^2 \phi_1^* = -(1/d_1)\eta_{2t}$. Substitution in (18a) gives:

$$\nabla^2 \psi_{1t} + \overline{\beta} \,\psi_{1x} = \frac{1}{d_1} (\overline{\beta} \,y \eta_{2t} + \partial_y \nabla^{-2} \eta_{2t}) \tag{20}$$

At the same time, the upper-layer equations (17b), (17d) have the same form as in the one-233 layer case, modulo a constant in front of the pressure gradient in the momentum equation, 234 and are treated analogously giving equation (9) for the upper-layer streamfunction, which has 235 asymptotic modon solutions. Equation (20) means that the equatorial modon in the upper 236 layer is accompanied by a linear Rossby-wave field in the lower layer, which is forced by 237 the pressure perturbations of the upper layer. Thus, the upper-layer modon is "leaky" and 238 loses energy through the generation of this Rossby-wave field, which means that its coherence 239 cannot be maintained for long time. 240

Equatorial Modons

241 2.2.4. Regime 2b

In this regime the deviation of the interface is much smaller than that of the free surface $\lambda_1 \sim \lambda, \lambda_2 \sim \lambda^2$, which means that we have an effectively barotropic situation. Indeed, the equations for upper- and lower-layer velocities become identical in the leading order in λ :

$$\begin{cases} \partial_{t} \mathbf{v}_{1} + \mathbf{v}_{1} \cdot \nabla \mathbf{v}_{1} + \overline{\beta} \, y \, \hat{\mathbf{z}} \wedge \mathbf{v}_{1} = -\frac{gH\lambda}{V^{2}} \nabla \overline{\eta}_{1}, \\ \partial_{t} \mathbf{v}_{2} + \mathbf{v}_{2} \cdot \nabla \mathbf{v}_{2} + \overline{\beta} \, y \, \hat{\mathbf{z}} \wedge \mathbf{v}_{2} = -\frac{gH\lambda}{V^{2}} \nabla \overline{\eta}_{1}, \\ \nabla \cdot \mathbf{v}_{1} = 0, \\ \nabla \cdot \mathbf{v}_{2} = 0. \end{cases}$$

$$(21)$$

Hence $\mathbf{v}_1 = \mathbf{v}_2$ and we get an asymptotic modon solution which is uniform through the whole fluid column, that is barotropic, and the situation is effectively equivalent to the one-layer case.

3. Numerical simulations with asymptotic modon initializations in one- and two- layer RSW models

251 3.1. Equatorial modon in different environments in one-layer model

Numerical simulations in the one-layer case were performed in a symmetric with respect to the 252 equator and wide enough domain, $9L_d \times 6.5L_d$, to minimize the boundary effects. Numerical 253 sponges allowing to absorb short inertia-gravity waves were implemented at the boundaries. 254 Here L_d is the equatorial deformation radius, $\sqrt{c/\beta}$, where $c = \sqrt{gH}$. We use the high-255 resolution, well-balanced finite-volume numerical scheme of Bouchut (2007). The dimensional 256 units of length and time in the numerical scheme were L_d and $1/(\beta L_d)$, respectively, which 257 corresponds to non-dimensional β equal to one. Dimensional values of q and β were fixed to be 258 those for the Earth, thus the mean depth H is a single parameter. If, for example, it is chosen 259 to be H = 10 km, which roughly corresponds to two thirds of the thickness of the tropical 260 troposphere, then the dimensional space and time units are $L_d \approx 3000 km, 1/\beta L_d \approx 3h$. Spatial 261 resolution varied from 600×600 to 1200×1200 , in order to check numerical convergence. No 262 external forcing nor dissipation or thermal relaxation were introduced. 263

²⁶⁴ 3.1.1. Adjustment of asymptotic modons, and formation of RSW modons

We first recall the results of the numerical simulations initialized with theoretical asymp-265 totic modon solutions, which were sketched in Rostami and Zeitlin (2019a), and give some 266 additional important details. The "dry" adiabatic RSW model is initialized with just a veloc-267 ity field acquired from the asymptotic modon solution (10), with $U \ll \sqrt{gH}$, and no pressure 268 perturbation. Nonlinear evolution of asymptotic modon leads to a process of relaxation pre-269 sented in Fig. 3, which consists in emission of short inertia-gravity waves and formation of a 270 dipolar coherent structure. Such "ageostrophic adjustment" of asymptotic modon to "real" 271 modon solutions of the full RSW equations was first observed in the simulations with RSW 272 model in the f-plane approximation in Ribstein et al. (2010). It lasts about one inertial period 273 $T = 1/(\beta L_d)$. After adjustment of pressure and velocity fields to twin cyclones on both sides 274 of the Equator, this coherent structure moves eastward, cf. Fig. 3. The phase speed of the 275 modon depends on the values of (a, U) used for initialization. We compared the evolution 276 starting from two sets of initial conditions corresponding to U = 0.1, U = 0.2 in (10), (11), 277 both with radius $a = 0.5L_d$. We should emphasize that there is a threshold $U \approx 0.07$ be-278

M. ROSTAMI & V. ZEITLIN



Figure 3. Initial "ageostrophic" adjustment of the asymptotic modon with parameters (a, p) = (0.5, 3.1623) corresponding to initialization with an asymptotic modon with U = 0.1, as seen in the thickness field.

low which a coherent dipole does not form from the initialization with respective asymptotic 279 modon. In both cases we have a formation of a coherent dipolar structure from the asymptotic 280 modon. However, while the weaker/slower dipole corresponding to U = 0.1 eventually slows 281 down and disaggregates at a distance $\approx 2 L_d$ from the initial location, the stronger one with 282 U = 0.2 continues to move eastward for much longer times. A comparison of the structure 283 of the dipoles arising from each initialization is presented in Fig. 4, their coherence following 284 from the scatter plot of potential vorticity vs Bernoulli function, cf. the left panel of Fig. 6. 285 The resulting RSW modons differ essentially from the parent asymptotic ones, as they has 286 a considerable thickness anomalies, cf. Fig. 4, with the respective values of non-dimensional 287 nonlinearity parameter ≈ 0.075 and ≈ 0.15 , respectively, but also a pronounced divergence 288 pattern, as follows from Fig. 5. 289

In spite of their coherence, we detected an enstrophy loss during the propagation of thus arising RSW modons, which diminishes with increasing resolution (Fig. 6). This loss of enstrophy because of numerical dissipation, which also affects the energy conservation, is not negligible for weaker/slower modon ($U = 0.1, a = 0.5L_d$), while it is reduced to just few % for stronger/faster modons ($U = 0.2, a = 0.5L_d$).

²⁹⁵ 3.1.2. Modons in adiabatic vs most-convective environment

We now switch on the effects of moisture within the simplest version of the diabatic moistconvective RSW (mcRSW) (Bouchut et al. 2009), in order to compare the behavior of equatorial modons in adiabatic and moist-convective environments. The equations of the model read:

$$\begin{cases} \partial_t \mathbf{v} + \mathbf{v} \cdot \nabla \mathbf{v} + \beta \, y \, \hat{\mathbf{z}} \wedge \mathbf{v} = -g \nabla h, \\ \partial_t h + \nabla \cdot (\mathbf{v}h) = -\gamma P, \\ \partial_t Q + \nabla \cdot (Q \mathbf{v}) = -P + E. \end{cases}$$
(22)

We introduced Q, a bulk amount of water vapor in the air column, with a condensation sink P and evaporation source E, which are parameterized as follows:

$$P = \frac{Q - Q^s}{\tau} \mathcal{H}(Q - Q^s), \quad E = \alpha |\mathbf{v}| (Q^s - Q) \mathcal{H}(Q^s - Q).$$
(23)

Here Q_s is a saturation moisture threshold, τ is relaxation time, α is a parameter regulating evaporation, γ is a parameter depending on background stratification, and $\mathcal{H}(...)$ denotes the Heaviside (step-)function. The parameterization of the condensation (23) is a bulk version of the Betts-Miller parameterization, and the parameterization of the evaporation is standard. As in the previous test simulations (Bouchut et al. 2009), we take a uniform initial moisture Equatorial Modons



Figure 4. Comparison of the structure of weaker (upper panel) and stronger (lower panel) modons, arising, respectively from initializations with U = 0.1 and U = 0.2. Left panel: thickness h at $T = 3[\beta L_d]^{-1}$. Middle panel: meridional section across the center of the modon. Right panel: zonal section of the equatorial modon across the center of each cyclone.

distribution Q_i close to saturation: $Q_i = Q_s - 0.01$ with $Q_s = 0.9$, a short relaxation time τ 309 equal to several time-steps of the numerical scheme, $\gamma = 1$, and $\alpha = \mathcal{O}(10^{-1})$. The values of 310 $Q_{i,s}$ are chosen to be the same as in (Bouchut et al. 2009) for benchmarking reasons; small 311 variations around them do not change qualitatively the results. A step-by-step comparison of 312 propagation of an equatorial modon in "dry" and moist-convective environments is presented 313 in Fig. 7. As follows from the Figure, the phase velocity of the modon in "dry" environment 314 is $\approx 0.025 L_d^2 \beta \approx 5m/s$, while in moist-convective environment it increases to $\approx 0.035 L_d^2 \beta \approx$ 315 12m/s, and the modon keeps its coherence. This is the effect of increase of the cyclonic 316 vorticity due to the moist convection, which is easy to understand within the mcRSW model, 317 cf. Lambaerts et al. (2011a). It is worth emphasizing that the phase velocity of a weak modon 318 is not constant because of enstrophy and energy losses due to numerical dissipation. This 319 is an indication that external friction forces are able to kill such modon. Nevertheless, the 320 convective enhancement and related increase of velocity allow the modon to run over $\approx 3L_d$, 321 which would be enough to pass over Indian Ocean, for example. Its scale, its propagation 322 speed, and associated convection patterns of this equatorial modon (cf. Rostami and Zeitlin 323 (2019a)) indicate a possible relevance to the MJO events. A specific feature introduced by the 324 steady zonal motion of equatorial modons is the mean zonal velocity which is induced in the 325 equatorial belt, and which can be of importance in the context of super-rotation of planetary 326 atmospheres. We present the zonally averaged velocity profile introduced by a modon, both 327 in adiabatic and moist-convective environments, in Fig. 8. 328

329 3.1.3. Interactions of equatorial modons with topography

We now investigate interactions of equatorial modons with topography, and changes in convective patterns introduced by this latter. For this, we consider a simplest configuration with an infinite zonally localized meridional ridge with a Gaussian profile of height, which is situated downstream from the advancing modon. The maximum height of the ridge is $b_{\text{max}} = 0.2H$ and it median width is equal to the radius of the core of the modon. The results are presented in Fig. 9. As seen in the Figure, the only effect of the ridge upon the

M. ROSTAMI & V. ZEITLIN



Figure 5. Snapshots at $T = 0, 10, 20, 30, 40, 50 [1/(\beta L_d)]$ after initial adjustment, from top to bottom, of h (left panel) and divergence (right panel) of the modon initialized with a = 0.5, U = 0.2.



Figure 6. Left panel: Scatter plot of Bernoulli function (mean value subtracted) in the co-moving frame vs potential vorticity (PV), which demonstrates the coherence of the modon. initialized with (a, p) = (0.5, 3.1623), corresponding to U = 0.1. Right panel: Sensitivity of the loss of enstrophy to numerical resolution.

geopotential and vorticity fields is a temporary lateral extension of the modon during the 336 passage over the ridge, which does not change neither its trajectory, nor coherence. Yet, the 337 zonal speed of the modon over topography abruptly decreases. Fig.10 shows evolution of 338 divergence and condensation patterns associated with the modon during the passage over the 339 ridge. Generation of inertia-gravity waves and of topographic waves, with related condensation 340 patterns is clearly seen. A weak signal corresponding to a reflected Rossby wave can be 341 identified on the left panel of Fig. 9, to the left of the ridge. Notice a typical condensation 342 pattern associated with the modon far from topography, which resembles that of the MJO 343 events. 344



Figure 7. Snapshots of pressure and velocity fields of the modon arising from an asymptotic modon initialization with U = 0.1 ((a, p) = (0.5, 3.1623)) in adiabatic (*left panel*) and moist-convective (*right panel*) environments.

M. ROSTAMI & V. ZEITLIN



Figure 8. Superposition of steady-state mean zonal velocity profile induced by the modon initialized with $a = 0.5, U = 0.2, \beta = 1$ in adiabatic (solid line) and diabatic (dashed line) environments.



Figure 9. Propagation of the equatorial modon in moist-convective environment over an idealized meridional ridge of Gaussian shape, as seen in the geopotential height (*left column*) and relative vorticity (*right column*) fields. Snapshots at T = 10, 20, 30, 40 [1/(βL_d)], from top to bottom. Maximum height of the ridge is 0.2*H*, and its median width is [-0.25]. Vertical dashed line represents the position of the center of the ridge. ($a = 0.5, U = 0.2, \alpha = 0.05$).

345 **3.2.** Two-layer Equatorial Modons

346 3.2.1. Adjustment of asymptotic modons, and existence of two-layer shallow-water modons in adiabatic environment

Following the strategy which proved to be fruitful in the one-layer model, and using the fact that the asymptotic modon solutions in Regimes 1 and 3 of section 2.2 are the same as in the one-layer model layerwise, we first initialize the simulations with "dry" two-layer equations (13) by the velocity field of the asymptotic modon solutions of section 2.1.2 in



Figure 10. Evolution of divergence (*left column*) and condensed water vapor (*right column*) fields during the passage of an equatorial modon over an idealized meridional ridge in moist-convective environment. Snapshots at T = 10, 20, 30, 40 $[1/(\beta L_d)]$, from top to bottom. Same conventions as in Fig. 9.

each layer, and no pressure perturbations, with a hope that via the process of ageostrophic 352 adjustment these configurations will end up in quasi-stationary states, like in the one-layer 353 case. Notice that such initialization fits all the regimes of section 2.2, as pressure anomalies of 354 the layers are initially identically zero. For simulations with two-layer model we are using the 355 numerical scheme of Bouchut and Zeitlin (2010), which was successfully tested in numerous 356 applications. In all simulations below the nonperturbed thicknesses of the layers are the same. 357 with the same total thickness as in the one-layer case, and the stratification parameter s is 358 equal to 1.1. The simulations were performed with initializations corresponding to asymptotic 359 modons of different sizes and intensities in respective layers. The overall conclusion following 360 from such experiments is that only the initializations with identical, or very close, asymptotic 361 modons in each layer, i.e. quasi-barotropic initializations, give rise to coherent slowly eastward-362 propagating structures, while essentially baroclinic initializations are eventually transformed 363 into equatorial Rossby-wave packets, although initially they form dipoles which may move 364 eastwards for some time. 365

A comparison of the evolution of strictly barotropic and weakly baroclinic initializations 366 with velocity fields corresponding, respectively, to the asymptotic modon of sec. 2.1.2 with 367 $U_1 = U_2 = 0.2$ in both layers , and with $U_1 = 0.2$ in the lower, and $U_2 = 0.1$ in the upper 368 layer is presented in Figure 11. As follows from the figure, the barotropic initialization leads 369 to a barotropic coherent dipole keeping its form and moving eastwards, while the dipole 370 produced by the weakly baroclinic initialization slows down and then engages in the reverse 371 westward motion losing its shape at the same time, that is gradually becoming a packet of 372 Rossby waves. The diagnostics of coherence with the scatter plots of Bernoulli function vs. 373 PV in the lower layer presented in Figure 12 confirm this conclusion. In the following we will 374 be calling stronger (weaker) baroclinic configurations those with larger (smaller) difference 375

M. ROSTAMI & V. ZEITLIN



Figure 11. "Dry" evolution of relative vorticity layer-wise starting from barotropic (first column: upper layer, second column: lower layer) and weakly baroclinic (third column: upper layer, fourth column: lower layer) initializations with velocity fields corresponding, respectively, to the asymptotic modon of sec. 2.1.2 with $U_1 = U_2 = 0.2$ in both layers, and with $U_1 = 0.2$ in the lower, and $U_2 = 0.1$ in the upper layer. Snapshots at $T = 0, 10, 20, 30, 50 [1/(\beta L_d)]$, from top to bottom.

between the amplitudes of initial parameters of asymptotic-modon initializations in the layers 376 $U_1 - U_2$. Our results show that strong baroclinic initializations do not produce coherent 377 long-living dipoles. A comparison of the evolution of strongly baroclinic initializations, with 378 asymptotic modon velocity parameter $U_2 = 0.2$ in the lower layer, with quiescent upper layer 379 $(U_1 = 0)$, and, inversely, with the same field in the upper layer, with quiescent lower layer 380 is presented in Figure 13, and shows a similar, modulo the layer swap, behavior displaying 381 a transformation of initial dipole into a wave-train. Analogous initialization with $U_1 = 0.1$, 382 i.e. smaller baroclinicity, produces a more coherent and long-leaving structure, see Figure 14 383 below. 384

385 3.2.2. 2-layer modons in moist-convective environment

To perform the two-layer simulations in moist-convective environment we work on the equatorial beta-plane with the simplest version of two-layer mcRSW with moist lower and dry upper layers, which was proposed in Lambaerts et al. (2011b) and ameliorated by including evaporation in Rostami and Zeitlin (2017). The equations of the model, in the same setting



Figure 12. Comparison of scatter plots of Bernoulli function (with subtracted mean value) vs PV in the lower layer of barotropic (*left panel*), and baroclinic (*right panel*) dipoles of Fig. 11 ($U_1 = 0.2, U_2 = 0.1$) at $T = 20 [1/(\beta L_d)]$, computed in a circle of radius 2a/3 with respect to the center of the dipole.



Figure 13. "Dry" evolution of relative vorticity in the lower layer (first column), and baroclinic velocity $v_1 - v_2$ and pressure $\eta_1 - \eta_2$ (second column) produced by essentially baroclinic initializations with the velocity parameter of the asymptotic modon of sec. 2.1.2 in the lower layer $U_1 = 0.2$, and quiescent upper layer: $U_2 = 0$. Same for initialization with asymptotic modon in the upper layer with $U_1 = 0, U_2 = 0.2$, respectively, the relative vorticity (third column), and baroclinic velocity and pressure (fourth column) in the upper layer. Snapshots at $T = 0, 15, 40, 60, 80 [1/(\beta L_d)]$, from top to bottom.

M. ROSTAMI & V. ZEITLIN

 $_{390}$ as in (13), read:

$$\begin{cases} \frac{D_1 \boldsymbol{v}_1}{Dt} + \beta y \, \hat{\mathbf{z}} \wedge \boldsymbol{v}_1 = -g \boldsymbol{\nabla} (h_1 + h_2), \\ \frac{D_2 \boldsymbol{v}_2}{Dt} + \beta y \, \hat{\mathbf{z}} \wedge \boldsymbol{v}_2 = -g \boldsymbol{\nabla} (h_1 + sh_2) + \frac{\boldsymbol{v}_1 - \boldsymbol{v}_2}{h_2} \beta P_1, \\ \frac{D_1 h_1}{Dt} + \boldsymbol{\nabla} \cdot (h_1 \boldsymbol{v}_1) = -\beta P_1, \\ \frac{D_2 h_2}{Dt} + \boldsymbol{\nabla} \cdot (h_2 \boldsymbol{v}_2) = +\beta P_1, \\ \frac{\partial_t Q_1 + \boldsymbol{\nabla} \cdot (Q_1 \boldsymbol{v}_1) = -P_1 + E. \end{cases}$$

$$(24)$$

Here $D_i/Dt = \partial_t + \boldsymbol{v}_i \cdot \boldsymbol{\nabla} \boldsymbol{v}_i$, i = 1, 2. The convective flux is linked to the water-vapour con-391 densation P_1 in the lower, humid layer, and produces a sink (source) in the mass conservation 392 equation in the lower (upper) layer, like in the one-layer mcRSW (22). However, now the total 393 mass is conserved. Due to the convective mass exchange the same flux leads to appearance of 394 the Rayleigh drag in the upper-layer momentum equation. Condensation and evaporation are 395 expressed in terms of the bulk moisture in the lower-layer column Q_1 in the same way as in 396 (23). The equations (22) arise from (24) in the limit of infinitely thick upper layer. The setup 397 is the same as in the "dry" simulations above. We take a uniform initial moisture distribution 398 in the lower layer close to saturation: $Q_1 = Q_s - 0.01$ with $Q_s = 0.8$, and choose $\alpha = 0.2$. 399

Before presenting the results of numerical experiments with two-layer mcRSW, let us recall 400 that condensation leads to enhancement of cyclonic vorticity in moist-convective environment, 401 as already mentioned. Hence, one can expect that, as in the one-layer case, the effects of 402 condensation make the dipoles arising from the adjustment of asymptotic modons more robust. 403 This is, indeed, what happens. In Figure 14 we show a comparison of "dry" and moist-404 convective evolution produced by the same as in Figure 13 initialization, but with a weaker 405 baroclinicity, with $U_1 = 0.1$, and quiescent upper layer. As follows from the Figure, the "dry" 406 modon in the lower layer keeps its coherence, while moving eastward for some time, but 407 then changes shape and engages in a westward motion becoming a looser dipole. The moist-408 convective evolution, with the same initial perturbation, maintains the eastward motion and 409 coherence of the initial modon in the lower layer for much longer times, with more intense 410 cyclonic vorticity compared to the "dry" case, although, eventually, the dipole is getting looser 411 and stops. 412

It would be tempting to interpret the westward-moving dipole arising in the "dry" scenario 413 in Figure 14 as a Rossby-wave soliton discovered by Boyd (1985) (which was also called modon) 414 and shown, theoretically and numerically, to arise in the equatorial adjustment of pressure 415 perturbations in (LeSommer et al. 2004). However, the Boyd's soliton is expected to arise at 416 much larger zonal to meridional aspect ratios, and was observed in numerical simulations in 417 a form of *anticuclonic* dipole. In fact, although the cyclonic dipole we observe maintains its 418 coherence for some time, it eventually loses it and becomes an equatorial Rossby-wave train 419 (not shown). 420

An interesting result is obtained with the barotropic initialization corresponding to the case of Figure 11 in moist-convective environment. It is presented in Figure 15, and shows that while the arising modon maintains it coherence for a long time, as follows from the vorticity snapshots (not shown), its baroclinic pressure pattern exhibits a quadrupolar structure, similar to that observed in the MJO events (Zhang 2005), with the condensation pattern also resembling that of MJO.



Figure 14. Comparison of "dry" (first and second columns) and moist-convective (third and fourth columns) evolution of relative vorticity in the lower layer (first and third panels), and baroclinic velocity $v_1 - v_2$ and pressure $\eta_1 - \eta_2$ (second and fourth panels) following an essentially baroclinic initialization with the velocity field in the lower layer corresponding to the asymptotic modon of sec. 2.1.2 with $U_1 = 0.1$, and quiescent upper layer: $U_2 = 0$. Snapshots at T = 0, 15, 40, 60, 80 $[1/(\beta L_d)]$, from top to bottom.



Figure 15. Snapshot at time at $T = 20 \left[\frac{1}{\beta L_d} \right]$ of baroclinic velocity $v_1 - v_2$ and pressure $\eta_1 - \eta_2$ (*left panel*) and condensation pattern (*right panel*) produced by barotropic initialization with the velocity field in both layers corresponding to the asymptotic modon of sec. 2.1.2 with U = 0.2.

427 4. Conclusion and Discussion

We, thus, established existence of nonlinear long-living, slow eastward-moving dipolar coherent structures, the modons, in the shallow-water dynamics in the equatorial beta-plane, both in one- and two-layer versions of the model, although robust modons are essentially barotropic in the latter case. These solutions of the fluid equations of motion at the Equator seem to be overlooked in the literature. The modons maintain their coherence in the presence

M. ROSTAMI & V. ZEITLIN

of moisture and are accompanied by a specific and persistent moist convection patterns, if 433 condensation and evaporation are included. The scales of these structures are conditioned 434 by the interpretation of the RSW model. At given g and β the actual scales depend on the 435 value of the unperturbed thickness H, or, equivalently, on the value of the corresponding 436 deformation radius L_d . A crucial observation which put us on the track of equatorial modons 437 is that Charney (1963) equation for equatorial balanced motions coincides with (9) modulo 438 parametric dependence on the vertical coordinate. Therefore, solutions of Charney's balance 439 equations in a form of asymptotic modons in the horizontal, with vertical modulation, do 440 exist, although it remains to prove that they survive in the full primitive equations. As we 441 showed, this is the case in one- and two-layer shallow-water models, provided the vertical 442 modulation is weak. 443

Coming back to a possible relation to MJO, we should emphasize that some features de-444 tected in numerical simulations we presented are indeed in resemblance with the MJO events. 445 These latter clearly display twin cyclonic structures in the lower layer (Rui and Wang 1990, 446 Yanai et al. 2000, Adames et al. 2014), and a quadrupolar structure of the first baroclinic 447 mode (Zhang 2005). An equivalent barotropic structure of MJO with a strong dipolar vortic-448 ity component was recently emphasized by Wang et al. (2019). Condensation/precipitation 449 patterns associated with equatorial modons resemble those of dipolar structures related to 450 MJO. Intensification and acceleration of the equatorial modons in moist-convective environ-451 ment is consistent with the observation that the regions of most active MJO are correlated 452 with the areas of large mean precipitation (Zhang and Dong 2004). 453

We should, however, say that, apparently, barotropic, or quasi-barotropic, modons alone 454 are not sufficient to explain the MJO events in all their complexity. It should not be forgotten 455 that these events do not maintain a fixed structure all the time, and have their own life-cycle, 456 an important part of which is related to interactions with Kelvin waves, e.g. (Haertel et al. 457 2014). A coupling of a baroclinic long-living modon-like structure in the moist-convective 458 environment, like that displayed in Figure 14, and a Kelvin wave, which was out of the scope 459 of the present study, can be shown to play a primordial role in the MJO dynamics (Rostami 460 and Zeitlin 2020). Yet, the coherent dipolar structures are certainly part of the MJO events, 461 and the increase of their eastward velocity with increase of intensity of the moist convection 462 is to be borne in mind in the context of global change and expected increase of temperature 463 of the oceans. 464

The existence of the slowly eastward-moving equatorial modons, and the scenarios of their 465 evolution, in particular in the presence of moist convection, should be not forgotten in data 466 analyses both on the Earth and on other planets. For example, the classical Gill's mechanism 467 (Gill 1980) was evoked in (Legarreta et al. 2016) with a support of "dry" RSW simulations, 468 in order to explain the so-called Y-shaped structures observed in the equatorial atmosphere 469 of Jupiter. The modifications of this mechanism with changes of parameters, and its ability 470 to generate equatorial modons, which were reported in (Rostami and Zeitlin 2019b), and also 471 the role of moist convection, which is known to be present in the Jovian atmosphere, as well 472 as the above-described scenario of transformation of eastward-moving coherent dipoles into 473 westward-moving and meridionally separating vortex pairs, should be kept in mind in this 474 context. 475

REFERENCES

476 References

- Adames, A. F., Patoux, J., Foster, R. C., 2014. The contribution of extratropical waves to the MJO wind field.
 J. Atmos. Sci. 71 (1), 155–176.
- Bouchut, F., 2007. Chapter 4: Efficient numerical finite volume schemes for shallow water models. In: Zeitlin,
 V. (Ed.), Nonlinear Dynamics of Rotating Shallow Water: Methods and Advances. Vol. 2 of Edited Series
 on Advances in Nonlinear Science and Complexity. Elsevier, pp. 189 256.
- Bouchut, F., Lambaerts, J., Lapeyre, G., Zeitlin, V., 2009. Fronts and nonlinear waves in a simplified shallowwater model of the atmosphere with moisture and convection. Phys. Fluids 21 (11), 116604.
- Bouchut, F., Zeitlin, V., 2010. A robust well-balanced scheme for multi-layer shallow water equations. Disc.
 Cont. Dyn. Sys. B 13 (4), 739–758.
- Boyd, J., 1985. Equatorial solitary waves. Part 3. Westward traveling modons. J. Phys. Oceanogr. 15, 46–54.
 Charney, J., 1963. A note on large-scale motions in the tropics. J. Atmos. Sci. 20, 607–609.
- Gill, A., 1980. Some simple solutions for heat induced tropical circulation. Q. J. Roy. Met. Soc. 106, 447–462.
 Gill, A., 1982. Atmosphere Ocean Dynamics. Academic Press.
- Haertel, P., Straub, K., Budsock, A., 2014. Transforming circumnavigating Kelvin waves that initiate and
 dissipate the Madden Julian Oscillation. Q. J. Roy. Met. Soc. 141, 1586–1602.
- Haertel, P., Straub, K., Fedorov, A., 2013. Lagrangian overturning and the Madden Julian oscillation. Q. J.
 Roy. Met. Soc. 140, 1344–1361.
- 494 James, I., 1994. Introduction to Circulating Atmospheres. Cambridge University Press.
- Kim, D., Sobel, A., Maloney, E., Frierson, D., Kang, I., 2011. A systematic relationship between intraseasonal
 variability and mean state bias in AGCM simulations. J. Climate 24, 5506–5520.
- Kizner, Z., Reznik, G., Fridman, B., Khvoles, R., McWilliams, J., 2008. Shallow-water modons on the f plane.
 J. Fluid Mech. 603, 305–329.
- Lahaye, N., Zeitlin, V., 2016. Understanding instabilities of tropical cyclones and their evolution with a moist
 convective rotating shallow water model. J. Atmos Sci. 73, 505–523.
- Lambaerts, J., Lapeyre, G., Zeitlin, V., 2011a. Moist versus dry barotropic instability in a shallow water model of the atmosphere with moist convection. J. Atmos. Sci. 68, 1234–1252.
- Lambaerts, J., Lapeyre, G., Zeitlin, V., Bouchut, F., 2011b. Simplified two-layer models of precipitating atmo sphere and their properties. Phys. Fluids 23, 046603.
- Larichev, V., Reznik, G., 1976. Two-dimensional solitary Rossby waves. Doklady USSR Acad. Sci. 231, 1077– 1080.
- Legarreta, J., Barrado-Izagirre, N., Garcia-Melendo, E., Sanchez-Lavega, A., Gomez-Forrellad, J., IOPW, 2016.
 A large active wave trapped in Jupiter's equator. A & A A15, A154.
- LeSommer, J., Reznik, G., Zeitlin, V., 2004. Nonlinear geostrophic adjustment of long-wave disturbances in
 the shallow-water model on the equatorial beta-plane. J. Fluid Mech. 515, 135–170.
- Madden, R., Julian, P., 1972. Description of global-scale circulation cells in the tropics with a 40-50 day period.
 J. Atmos. Sci. 29, 1109–1123.
- 513 Majda, A. J., Stechman, S. N., 2009. The skeleton of tropical intraseasonal oscillations. PNAS 106, 8417–8422.
- Matsuno, T., 1966. Quasi-geostrophic motions in the equatorial area. J. Met. Soc. Japan 44, 25–43.
- Ribstein, B., Gula, J., Zeitlin, V., 2010. (A)geostrophic adjustment of dipolar perturbations, formation of
 coherent structures and their properties, as follows from high-resolution numerical simulations with rotating
 shallow water model. Phys. Fluids 22, 116603.
- Rostami, M., Zeitlin, V., 2017. Influence of condensation and latent heat release upon barotropic and baroclinic
 instabilities of vortices in a rotating shallow water f-plane model. Geophys. Astrophys. Fluid Dyn. 111 (1),
 1-31.
- Rostami, M., Zeitlin, V., 2018. Improved moist-convective rotating shallow water model and its application to
 instabilities of hurricane-like vortices. Q. J. Roy. Met. Soc. 144, 1450–1462.
- Rostami, M., Zeitlin, V., 2019a. Eastward-moving convection-enhanced modons in shallow water in the equatorial tangent plane. Phys. Fluids 31, 021701.
- Rostami, M., Zeitlin, V., 2019b. Geostrophic adjustment on the equatorial beta-plane revisited. Phys. Fluids
 31, 081702.
- Rostami, M., Zeitlin, V., 2020. Can geostrophic adjustment of baroclinic disturbances in tropical atmosphere
 explain MJO events? Q. J. Roy. Met. Soc. https://doi.org/10.1002/qj.3884.
- Rui, H., Wang, B., 1990. Development characteristics and dynamical structure of tropical intraseasonal con vection anomalies. J. Atmos. Sci. 47, 357–379.
- Sanchez-Lavega, A., Lebonnois, S., Imamura, T., Read, P., Luz, D., 2017. The atmospheric dynamics of Venus.
 Space Sci. Rev. 212, 1541–1616.
- Showman, A. P., Dowling, T., 2000. Nonlinear simulations of Jupiter's 5-micron hot spots. Science 289, 1737–
 1740.
- Sobel, A., Maloney, E., 2012. An idealized semi-empirical framework for modeling Madden-Julian oscillation.
 J. Atmos. Sci. 69, 1691–1705.
- Straub, K., Kiladis, G. N., 2002. Observations of a convectively coupled Kelvin wave in the Eastern pacific
 ITCZ. J. Atmos. Sci. 59, 30–52.
- Tribbia, J., 1984. Modons in spherical geometry. Geophys. Astrophys. Fluid Dyn. 30, 131–168.
- Verkley, W., 1984. The construction of barotropic modons on the sphere. J. Atmos. Sci. 41, 2492–2504.

0	ົ
4	4

REFERENCES

- Wang, D., Yano, J.-I., Lin, Y., 2019. Madden Julian oscillations seen in the upper-Troposphere vorticity field:
 interactions with Rossby wave trains. J. Atmos. Sci. 76, 1785–1807.
- Wheeler, M., Kiladis, G., 1999. Convectively coupled equatorial waves: analysis of clouds and temperature in
 the wavenumberfrequency domain. J. Atmos. Sci. 56, 374–399.
- Yanai, M., Chen, B., Tung, W.-W., 2000. The Madden Julian Oscillation observed during the TOGA COARE
 IOP: Global View. J. Atmos. Sci. 57 (15), 2374–2396.
- Yano, J.-I., Bonazzola, M., 2009. Scale analysis for large-scale tropical atmospheric dynamics. J. Atmos. Sci.
 66, 159–172.
- Yano, J.-I., Mulet, S., Bonazzola, M., 2009. Tropical large-scale circulations: asymptotically non-divergent?
 Tellus 61A, 417–427.
- Yano, J.-I., Tribbia, J., 2017. Tropical atmospheric Madden–Julian Oscillation: a strongly nonlinear free solitary
 Rossby wave? J. Atmos. Sci. 74, 3473–3489.
- Zeitlin, V., 2018. Geophysical Fluid Dynamics: Understanding (almost) Everything with Rotating Shallow
 Water Models. Oxford University Press.
- 555 Zhang, C., 2005. Madden-Julian oscillation. Rev. Geophys. 43, RG2003.
- ⁵⁵⁶ Zhang, C., Dong, M., 2004. Seasonality in the Madden Julian oscillation. J. Climate 17 (16), 3169–3180.